

Seizures

Seizures are caused by abnormal electrical firing in the brain. In some cases, the seizures are caused by an identifiable problem inside the brain (i.e. cancer, infection, inflammation, congenital brain malformation). Animals with a known cause are said to have symptomatic epilepsy.

What causes a seizure?

Seizures are caused by abnormal electrical firing in the brain.

In some cases, the seizures are caused by an identifiable problem inside the brain (i.e. cancer, infection, inflammation, congenital brain malformation). Animals with a known cause are said to have symptomatic epilepsy. These dogs often have other neurologic signs (i.e. changes in behavior, difficulty walking, circling). In some animals with symptomatic epilepsy, the seizures stop with anti-seizure treatment and/or treatment of the underlying disease process.

In animals with primary or idiopathic epilepsy, there is no identifiable abnormality inside the brain to cause the seizures. This form of epilepsy is believed to be genetic or inherited. Primary epilepsy occurs in approximately 5% of dogs. Common breeds include German Shepherds, German Shorthaired Pointers, Labrador Retrievers, Golden Retrievers, Border Collies, Shetland Sheepdogs, and English Springer Spaniels.

Seizures in dogs with primary epilepsy typically begin at 6 months – 5 years of age. Juvenile or late adult onset epilepsy occurs in some dogs but is very rare.

How do I know whether my dog is having a seizure?

There are many different types of seizures. The most common is called a generalized or “grand mal” seizure and is characterized by violent movement of all four limbs while lying down. Chomping or tightening of the jaw may occur. Animals are typically unresponsive during a seizure and commonly urinate, defecate, and salivate.

Some animals have partial or focal seizures, which may manifest as twitching of the face, chomping of the jaw, movement of a limb, or moments of unresponsiveness, confusion, or staring into space.

Vienna

The Hope Center
140 Park St, SE
Vienna, VA 22180
p 703.242.6000
f 703.242.8358

Leesburg

The LifeCentre
165 Fort Evans Rd, NE
Leesburg, VA 20176
p 703.771.2100
f 703.777.9968

Winchester

Valley Veterinary Emergency
and Referral Clinic
210 Costello Dr
Winchester, VA 22602
p 540.450.0177
f 540.662.7870

Woodbridge

Woodbridge Animal Hospital
2703 Caton Hill Rd
Woodbridge, VA 22191
p 703.242.6000
f 703.242.8358

Rockville

Metropolitan Emergency
Animal Clinic
12106 Nebel St
Rockville, MD 20852
p 301.816.0177
f 301.816.0179



Many animals with primary epilepsy have a classic pattern to their seizures. They may occur at a certain time interval (i.e. every 3 weeks, every 4 months), a certain time of day (commonly in the early morning) or in association with a particular stressful activity or event (i.e. guests in the home, going to the groomer, owner out of town).

Some dogs have a pattern of "cluster" seizures, characterized by 2 or more seizures within a defined period (i.e. 6 seizures in 48 hours). Dogs with cluster seizures are often more difficult to control with the standard anti-seizure medications.

Many dogs have a period before the seizure (called the pre-ictal period) where they demonstrate particular behavior patterns (i.e. neediness, anxiety, isolation, sedation), which may last hours to days. Similarly, many dogs have a classic period after the seizure (called the post-ictal period) where they exhibit unusual behaviors (i.e. seeming blind, unsteady when walking, excessively hungry or thirsty, excessively sedate), which may also last hours to days.

Some conditions can look similar to seizures including fainting episodes caused by heart or respiratory disease, episodes of loss of balance (called vestibular episodes or events), behavioral conditions (i.e. fly biting, tail chasing), and head bobbing syndromes (occur commonly in English Bulldogs, Doberman Pinschers, and Labrador Retrievers).

Are seizures harmful for my pet?

Although extremely stressful to witness, short seizures (<1 minute-2 minutes) are typically not harmful. Seizure activity becomes dangerous when patients enter a stage called status epilepticus, which is defined as a seizure lasting longer than 5 minutes or having multiple seizures without regaining consciousness in between. Possible consequences of status epilepticus include swelling in the brain, extremely high body temperature, and systemic complications including abnormalities in blood clotting.

Why and when is it important to treat seizures?

We recommend treating epilepsy when dogs have frequent seizures (i.e. greater than every 1-2 months) or severe seizures (lasting longer than 3-5 minutes). In untreated animals, seizures may become more severe or more frequent over time.

What are the most common anti-seizure medications and their side effects?

The most common anti-seizure medications used in veterinary

Vienna

The Hope Center
140 Park St, SE
Vienna, VA 22180
p 703.242.6000
f 703.242.8358

Leesburg

The LifeCentre
165 Fort Evans Rd, NE
Leesburg, VA 20176
p 703.771.2100
f 703.777.9968

Winchester

Valley Veterinary Emergency
and Referral Clinic
210 Costello Dr
Winchester, VA 22602
p 540.450.0177
f 540.662.7870

Woodbridge

Woodbridge Animal Hospital
2703 Caton Hill Rd
Woodbridge, VA 22191
p 703.242.6000
f 703.242.8358

Rockville

Metropolitan Emergency
Animal Clinic
12106 Nebel St
Rockville, MD 20852
p 301.816.0177
f 301.816.0179

medicine are Phenobarbital and Potassium Bromide. They have the advantage of being inexpensive.

Phenobarbital

Dosages: 15 mg, 30 mg, 60 mg; 1/4 grain (16.2 mg), 1/2 grain (32.4 mg), 1 grain (64.8 mg) tablets

Dosing schedule: Typically twice daily (every 12 hours); important not to miss a dose; stopping suddenly can cause seizures

Time to take effect: reaches full effect approximately 2 weeks after starting the medication

Species: Okay for use in dogs and cats

Common side effects:

Sedation, incoordination when walking – common in the first 1-2 weeks after starting Phenobarbital or increasing the dose; may persist in some dogs especially if on high doses or in dogs sensitive to these side effects

Increased drinking, urination, and appetite – typically persist while an animal is on the medication

Increased liver metabolism resulting in increased liver enzyme values – this is normal and not an indication of liver damage

Very rare side effects:

Reversible liver damage – evidenced by decreased production of proteins by the liver; can be monitored with routine bloodwork every 6-12 months

Reversible bone marrow damage; can be monitored with routine bloodwork every 6-12 months

Behavior changes (i.e. aggression)

Potassium bromide

Doses: 500 mg, 1000 mg capsules; 200 mg/mL or 250 mg/mL solutions

Dosing schedule: Once daily but can be given twice daily; missing dose is less hazardous due to long half life in blood stream

Timing to take effect: Takes 2-3 months to take effect when started on maintenance dose; therefore most dogs are given a loading dose to make it take effect sooner

Species: Okay for use in dogs only

Common side effects:

Sedation, incoordination when walking – common in the first 1-2 weeks after starting potassium bromide or increasing the dose; may persist in some dogs especially if on high doses or in dogs sensitive to these side effects

Increased drinking, urination, and appetite – typically persist while a dog is on the medication

Gastrointestinal upset – common if given without food or if using the capsule rather than the liquid form Fatal lung disease in



Vienna

The Hope Center
140 Park St, SE
Vienna, VA 22180
p 703.242.6000
f 703.242.8358

Leesburg

The LifeCentre
165 Fort Evans Rd, NE
Leesburg, VA 20176
p 703.771.2100
f 703.777.9968

Winchester

Valley Veterinary Emergency
and Referral Clinic
210 Costello Dr
Winchester, VA 22602
p 540.450.0177
f 540.662.7870

Woodbridge

Woodbridge Animal Hospital
2703 Caton Hill Rd
Woodbridge, VA 22191
p 703.242.6000
f 703.242.8358

Rockville

Metropolitan Emergency
Animal Clinic
12106 Nebel St
Rockville, MD 20852
p 301.816.0177
f 301.816.0179

CATS

Rare side effects:

Inflammation of the pancreas – may require hospitalization and warrants discontinuation of the medication in some dogs

Other medications (i.e. Keppra, Zonisamide) are often used in dogs with severe side effects to Phenobarbital and Potassium Bromide or in dogs with a high seizure frequency despite receiving Phenobarbital and Potassium Bromide.

Is it okay to leave my pet alone? What happens if a seizure occurs when I'm not home?

Many owners fear leaving their pets at home and feel that they need constant monitoring. Although it is natural to be concerned, in reality we cannot be with our pets 24 hours/day. Most animals have short seizures, and they will recover on their own. Signs that your pet has seized when you are gone include finding urine, feces, or drool in the house or finding our pet seeming dazed or disoriented (in the post-ictal phase) when you come home.



Vienna

The Hope Center
140 Park St, SE
Vienna, VA 22180
p 703.242.6000
f 703.242.8358

Leesburg

The LifeCentre
165 Fort Evans Rd, NE
Leesburg, VA 20176
p 703.771.2100
f 703.777.9968

Winchester

Valley Veterinary Emergency
and Referral Clinic
210 Costello Dr
Winchester, VA 22602
p 540.450.0177
f 540.662.7870

Woodbridge

Woodbridge Animal Hospital
2703 Caton Hill Rd
Woodbridge, VA 22191
p 703.242.6000
f 703.242.8358

Rockville

Metropolitan Emergency
Animal Clinic
12106 Nebel St
Rockville, MD 20852
p 301.816.0177
f 301.816.0179